For Immediate Release Lawson Health Research Institute and Western University April 25, 2019

Researchers verify new method of HIV transmission among injection drug users and effective prevention technique

"Cook your Wash" campaign reduces risk of HIV transmission

LONDON, ON – New studies from Lawson Health Research Institute and Western University have found for the first time that HIV can be transmitted through the sharing of equipment used to prepare drugs before injection and that a simple intervention – heating the equipment with a cigarette lighter for 10 seconds – can destroy the HIV virus, preventing that transmission. The findings, used to inform a public health campaign called 'Cook Your Wash,' have helped reduce rates of HIV transmission in London, Ontario.

The two studies published in *Journal of Acquired Immune Deficiency Syndrome (JAIDS)* were initiated in 2016 to address a public health emergency in London, Ontario when HIV rates amongst injection drug users more than doubled.

"This outbreak occurred despite London having Canada's largest per capita sterile needle and syringe distribution program, a strong opiate substitution therapy program and a multidisciplinary HIV clinic," says Dr. Michael Silverman, Lawson Associate Scientist and Chair/Chief of Infectious Diseases at the Schulich School of Medicine & Dentistry, Western University, London Health Sciences Centre and St. Joseph's Health Care London. "We knew there must be a novel method of HIV transmission at play."

From August 2016 to June 2017, the research team interviewed 119 injection drug users to understand their injection behaviours and risk for HIV. They discovered that those who shared equipment used to prepare drugs for injection were 22 times more likely to contract HIV than those who did not, despite not sharing needles or syringes.

The equipment includes a metal 'cooker' used to dissolve drugs in water and a filter used to draw the mixture, known as 'the wash,' into the syringe. Injection drug users reported reusing the equipment when consuming controlled-release hydromorphone, one of the most commonly injected opioids.

"Controlled-release hydromorphone is expensive and difficult to dissolve. After the first wash, large amounts of the drug remain in the equipment which is then saved, shared or sold for future use," explains Dr. Sharon Koivu, Associate Scientist at Lawson and Associate Professor at Schulich Medicine & Dentistry. "While people know not to share needles, some use their own needle multiple times allowing for contamination of the equipment."

The team took their findings back to the research laboratory. They confirmed that, on average, 45 per cent of the drug remains in the equipment after the first wash. They not only confirmed the HIV virus can be transmitted between needles, cookers and filters, but also discovered that controlled-release hydromorphone has properties that promote survival of the virus.

"The slow release properties in the drug can unfortunately stabilize the HIV virus," says Eric Arts, PhD, Chair of the Department of Microbiology and Immunology at Schulich Medicine & Dentistry. "For the first time, we were able to demonstrate that sharing equipment could lead to the type of HIV outbreak we observed in the community."

The team discovered that when the cooker is heated with a cigarette lighter for approximately 10 seconds, or until the wash bubbles, the virus is destroyed. They termed the technique 'cook your wash.'

They also confirmed that heating the cooker did not impact drug concentration. "We had to make sure that cooking your wash would not change the amount of drug being drawn into the syringe," notes Dr. Silverman. "If too much drug was released, it could lead to overdoses. If any drug was burnt off or lost, the intervention would not be accepted by persons who inject drugs."

Partnering with local community organizations like the Middlesex-London Health Unit and Regional HIV/AIDS Connection, 'Cook Your Wash' was launched as a public health campaign.

"The 'Cook Your Wash' campaign is one of the most exciting things to happen in our community," says Dr. Koivu. "We learned from persons who inject drugs, took that information to the lab to develop a solution and then brought that solution back to the community in record time."

Dr. Silverman adds that local rates of new HIV cases fell dramatically after the introduction of the campaign. "It wasn't the sole reason for the reduction in HIV rates as other interventions were also introduced, but the timing suggests it was part of the solution."

The studies build on previous research that shows sharing equipment can lead to the transmission of hepatitis C and controlled-release hydromorphone can promote the survival of bacteria that can cause endocarditis, which is an infection of the heart valves.

The team hopes this research can be used to inform understanding and interventions in other centres facing HIV epidemics amongst persons who inject drugs. "We hope our findings can be used to reduce the incidence of HIV transmissions even further and that, one day, society will be HIV free," says Dr. Silverman.

Video of the researchers discussing their work can be found here: <u>https://youtu.be/VQK3SQFzxTQ</u>.

-30-

MEDIA CONTACT: Robert DeLaet, Communications and External Relations, Lawson Health Research Institute, t. 519.685.8500 ext. 75664, c. 519.619.3872, <u>robert.delaet@lawsonresearch.com</u>

ABOUT LAWSON HEALTH RESEARCH INSTITUTE

As the research institute of London Health Sciences Centre and St. Joseph's Health Care London, and working in partnership with Western University, Lawson Health Research Institute is committed to furthering scientific knowledge to advance health care around the world.

ABOUT WESTERN

Western University delivers an academic experience second to none. Since 1878, The Western Experience has combined academic excellence with life-long opportunities for intellectual, social and cultural growth in order to better serve our communities. Our research excellence expands knowledge and drives discovery with real-world application. Western attracts individuals with a broad worldview, seeking to study, influence and lead in the international community.

ABOUT THE SCHULICH SCHOOL OF MEDICINE & DENTISTRY

The Schulich School of Medicine & Dentistry at Western University is one of Canada's preeminent medical and dental schools. Established in 1881, it was one of the founding schools of Western University and is known for being the birthplace of family medicine in Canada. For more than 130 years, the School has demonstrated a commitment to academic excellence and a passion for scientific discovery.